

EFFECT OF EXPERIMENTAL LIVER FLUKE (*Fasciola gigantica*) INFECTION ON LIVE WEIGHT CHANGES AND GROSS PATHOLOGY OF SAVANNAH BROWN GOATS RAISED UNDER SEMI-INTENSIVE SYSTEM.

Adama, J.Y.

Department of Animal Production,
Federal University of Technology, Minna
Amend- adama.live@yahoo.com
FUT email – john.adama@futminna.edu.ng

Received 13th July, 2016, Accepted 26th September, 2016

Abstract

The aim of this study was to determine the effect of experimental *Fasciola gigantica* infection on live weight changes and possible gross pathological lesions in Savannah Brown goats raised semi-intensively. Fifteen Savannah Brown goats were used for this experiment and the experimental design was Randomised Complete Block Design (RCBD). Two weeks pre-infection data was collected, and they were randomly assigned to three treatment groups with treatment 1 (infected animals without supplementation) and treatment 2 (infected with supplementation) and treatment 3 (control), respectively. The infected animals were each given 300 metacercariae orally, and all the animals were allowed to graze on natural pasture for about five (5) hours daily while they were fed with beans husk as supplement and fresh clean water was also provided *adlibitum* throughout the experiment. Significant differences ($P < 0.05$) were observed in the live weight of the infected groups as from week 8 to week 12 post-infection when compared to the respective values of the controls. The mean live weights of the infected groups from week 8 to week 12 were; T1 (6.5 ± 5.5 , 6.1 ± 5.1 , 5.8 ± 5.1 , 4.5 ± 5.5 and 4.3 ± 5.3), T2 (10.43 ± 1.80 , 9.8 ± 5.0 , 7.9 ± 5.0 , 7.4 ± 4.5 and 4.8 ± 5.5) compared to those for the control group (16.0 ± 4.7 , 16.0 ± 4.7 , 16.0 ± 4.7 , 17.0 ± 4.7 , 17.0 ± 4.7). The gross pathological lesions observed in dead goat from the infected group without supplementation was distended gall bladder, haemorrhagic liver as well as areas of necrosis. It was therefore, recommended that supplementary feeding during dry season periods of the year be practiced by livestock farmers within the study area in order to minimize the effect of disease conditions among ruminant livestock.

Keywords: *Fasciola gigantica*, Experimental infection, gross pathological lesions.

Introduction

Goat production is a specialized business since its operation requires some skills like good feeding, proper health and housing management, a deviation from which may lead to a variety of problems such as disease outbreak, parasitic infestation and poor output (Lasisi, 2010).

The population of goats in Nigeria numbered about 35 million (Adebawale, 2012), but the quantity of meat derived from these animals is below demand and this is partly due to the effects of parasitic diseases, such as helminthosis which causes a major health problem that lead to unprofitable production of small ruminants in the country.

A significant proportion of our ruminant livestock in Nigeria and in the tropics are reared under transhumance system with supplementary feeding, resulting in low productivity and high pre-weaning mortality (Adama *et al.*, 2011). Similarly acute shortage of

feeds during the dry season remains a common occurrence, this compel these animals to graze in areas that are often heavily infested with the potential intermediate hosts of liver fluke (Traor, 1986; Adama *et al.*; 2011), leading to disease outbreak such as fasciolosis caused by *Fasciola gigantica*.

Studies across the country have shown that helminthosis are known to be widespread in Nigeria (Biu *et al.*, 2009, Jatau, 2011) and that helminth parasites are by far the most serious causes of production losses in farm ruminants and that they are indisputably the cause of serious production losses to ruminants in Sub-Saharan Africa and worldwide (Ng'ang'a *et al.* 2004; Rahamn, 2004). This experiment is therefore designed to determine the live weight changes and gross pathological lesions observed in *Fasciola gigantica* infected Savannah Brown goats under semi-intensive of management system.

Materials and Methods

Study Area

This study was carried out at the Research farm of the Department of Animal Production, Federal University of Technology, Minna in Bosso Local Government Area of Niger State located in Southern Guinea Savanna zone of Nigeria. Minna has a land mass of 28.55sq. Km, and lies between longitude 6°29' of north, the average temperature range is between 18 to 39°C with an average monthly rainfall of 1200mm (NSADP, 1995).

Source of Experimental Animals and Feed

Fifteen (15) Savannah Brown goats of mixed sex, aged between 6 months to 1 year old and weighing between 7 -10kg were obtained in Beji market, Bosso Local Government Area of Niger State. Similarly, the feed supplement, 1,000grams of Beans Husk (*Vigna sinensis*) was also purchased in the market.

Pre-infection Management

The animals used for the experiment were properly screened against parasitic infections and other blood parasites through the use of Ivomec injection at 0.5ml/10kg subcutaneously, and long acting Terramycin injection (TLA) at 1ml/10kg body weight through deep intramuscular muscle. After the treatment, they were further screened to ensure that they were free of any possible infection. Two weeks pre-infection live weights were obtained in this regard.

Experimental Animals

The experiment was arranged in Randomized Complete Block Design (RCBD). Fifteen Savannah Brown goats were used in the experiment. The animals in two groups were each given 500g of beans husk daily for 3 months while the control group was not given any supplement, water was given *adlibitum* after grazing for the day. There was equal representation of the different ages and sex in the three groups. Baseline pre-infection data were collected and the goats were randomly assigned to three treatment groups.

Infective Materials

Viable *Fasciola gigantica* metacercariae were obtained from the Department of Parasitology Laboratory, Faculty of Veterinary Medicine, A.B.U Zaria. They were packed in ice bag and later stored at 4°C in the refrigerator to ascertain their viability till the day of usage.

Animal Infection

The fifteen (15) animals were randomly divided into three groups of five animals each, representing infected with supplement (T₁), infected without supplement (T₂) and control with supplement (T₃). Each of the animals in the infected group was inoculated orally with 300 *Fasciola gigantica* metacercariae according to the procedure described by Adama *et al.* (2011). The experiment is such that all the infected animals were treated after the experiment in order not to bridge ethical values.

Management of Experimental Animals

All animals were managed under semi-intensive system. Animals were let out to graze freely on natural graze land between 8:00 am to 3:00 pm and they were returned back to their pen and provided feed supplement, bean husks and clean fresh water was also given. The entire study lasted for twelve (12) weeks.

Data Collection

Live weight changes were monitored weekly for a period of ten (10) weeks after (2) weeks pre-infection monitoring using a weighing scale, manufactured from Hana Company with T number 005567. Gross pathological lesions were also monitored in some infected animals through pictorial views.

Data Analysis

The data collected were subjected to Analysis of Variance (Anova) using statistical package (SAS, 2008). Means were separated by the least significant Difference method (Steel and Torrie, 1980).

Results and Discussion

The result of the weight changes observed in the experiment are presented in Table 1 which indicates that there was no significant difference ($P > 0.05$) in the body weight changes up to a period of seven (7) weeks. However, significant differences were obtained from week 8 to 12 between the infected treatment without supplement (T₁) and the infected treatment with supplement (T₂) as well as the controls respectively. From 8 -12th week of the experimental period, the weight changes obtained in the control group differs significantly ($P < 0.05$) from those of (T₁) and (T₂) respectively. Similarly, during the same period the weight changes obtained in (T₁) differs significantly from 8 to 11th week of the experiment with no significant ($P > 0.05$) difference observed at the 12th week of the experiment between the two treatments.

Table 1: Mean body weight of Savanna Brown goats infected with 300 *Fasciola gigantica* metacercariae with or without supplementation with beans husks (Corps pod).

Weeks	T ₁	T ₂	T ₃	SEM	LS
1	10.8±3.6	11.0±1.8	12.0±4.7	1.0	NS
2	10.8±3.6	11.0±1.8	12.0±4.7	1.0	NS
3	10.8±3.6	10.0±1.8	13.0±4.7	1.1	NS
4	10.8±3.6	10.0±1.8	13.0±4.7	1.1	NS
5	11.3±4.1	11.45±2.4	14.0±4.7	1.2	NS
6	11.0±3.9	11.44±2.6	14.0±4.7	1.2	NS
7	11.0±3.9	11.32±2.6	15.0±4.7	1.3	NS
8	6.5±5.5 ^b	10.43±1.80 ^{bc}	16±4.7 ^a	1.4	*
9	6.1±5.1 ^b	9.8±5.0 ^{bc}	16.0±4.7 ^a	1.7	*
10	5.8±5.1 ^b	7.9±5.0 ^{bc}	16.0±4.7 ^a	1.8	*
11	4.5±5.5 ^b	7.4±4.5 ^{bc}	17.0±4.7 ^a	2.0	*
12	4.3±5.3 ^b	4.8±5.5 ^b	17.0±4.7 ^a	2.0	*

T₁ =infected without supplement, T₂ = Infected with supplement, T₃ = Control,
 abc = means on the same row with different superscript are significantly different(P<0.05).
 NS = No significant difference (P>0.05), L_s = level of significance, Sem = Standard error of mean

The gross pathological lesions of the liver of a goat in the infected group without supplement (T₁), shows a distended gall bladder with haemorrhagic liver (plate1). A further opening up of the same liver (plate11) shows gross pathological lesions of haemorrhages and areas of necrosis.

A picture of a *Fasciola gigantica* infected goats in infected group without Supplement (T₁) at week 7 post infection is shown in plate III. Similarly recovered immaturred worms from the liver of a goat without supplementation (T₁) is shown in plate IV.

The significant (P<0.05) weight losses recorded in this study from the two infected groups(T1 and T2) when compared to the control (T3) as from week 8 to 12 as a result of experimental *Fasciola gigantica* infection is consistent with earlier reports observed in small ruminants (Radostits *et al*; 2000; Egual *et al*; 2009 and Adama *et al*; 2011) that trematodes cause serious economic losses mostly through reduction in weight and productivity as well as increased mortality. The higher weight gain observed in T₃ (control group) confirmed earlier report (Hussein *et al*; 1983) that goats gain more weight and utilize feed more efficiently under supplementary feeding, particularly under stall fed conditions.

Post mortem examination of the dead animal shows haemorrhagic liver, with areas of necrosis due to extensive liver damage as a result of migration through the liver tissues, this clearly agrees with the findings of (Egual *et al*;

2009), the authors worked on experimental infection of Ethiopian sheep breeds using *Fasciola hepatica* and recovered several immature flukes from the liver parenchyma showing extensive liver damage. The pathology observed was further confirmed by the recovery of immature flukes from the liver parenchyma which might have caused some levels of damage to the liver cells.

Conclusion

It was concluded that the supplemented infected group showed higher resistance to *Fasciola gigantica* infection as compared to the non-supplemented infected group. Similarly, gross pathological observations revealed greater damage to liver tissues in non-supplemented infected animal that died at week 10 post-infection. It is recommended that supplemental feeding particularly during dry season periods of the year be given to small ruminants in order to minimize the effect of disease conditions



Plate 1: Distended gall bladder as a result of *Fasciola gigantica* infection on the goat that died at 10th week post-infection. Note: distended gall bladder (G).




Plate II: Haemorrhagic liver showing the area of necrosis (N) on the infected liver at 10th week Post-infection. Note: Areas of haemorrhage (H).



Plate III: A picture of an emaciated goat without supplementation (T₂) as a result of *Fasciolagigantica* infection at 7th week post-infection.



Plate IV: Recovered immature worms from the liver of a goat without supplementation (T₂) at 10th week post-infection

References

- Adama, J.Y., Ajanusi, O.J., Chiezey, N., Lawal, A. (2011). Haematological responses of Yankasa sheep to experimental *Fasciola gigantica* infection in Zaria, Nigeria. *Agriculture and Biology Journal of North America*. Volume 2(8): 1232-1238.
- Adebowale L.O.A (2012). Dynamics of Ruminant Livestock Management in the context of the Nigerian Agricultural System. In livestock production. Khalid, J(ed). In Tech. chapters publishers.
- Biu, A. A., Maimunatu, A., Salamatu, A. F. and Agbadu, E. T. (2009). A faecal survey of gastrointestinal parasites of ruminants on the University of Maiduguri Research Farm. *International Journal of Biomedical and Health Sciences* Vol. 5, No 4. December 31, 2009.
- Egualle, T., C.A. Mekonnen, and H. Chaka (2009). Evaluation of variation in susceptibility of Ethiopian sheep breeds to experimental infection with *Fasciola hepatica* response volume, 3, doi:10.1016/Journal of Small Ruminant Research. 2008 12-17.
- FAO (2008). Food and Agricultural Organization Production Year Book.
- Hussain, M.Z., Naidu, R., Tavuki and Sigh, R. (1983). Goat production and development in Fiji wild Animal Response, 48: 25-35.
- Jatau, I. D., Abdulganiyu, A., Lawal, A. I., Okubanjo, O. O. and Yusuf, K. H. (2011). Gastrointestinal and Haemoparasitism of sheep and goats at slaughter in Kano, Northern-Nigeria.
- Millar, D. G., Griffith, S., Algae, E. K, (1982): Activity and starting of glycolytic Enzymes in the presence of ethanol Biotechnol 601-606.
- Moulin, G., Booze H., and Galzy, (1984). Inhabitation of alcoholic fermentation Biotechnol. Bioeng 25365-382
- Saliu B. K. and Sani A. (2012) Bioethanol potentials of corn cob hydrolysed using cellulases of *Aspergillus niger* and *Penicillium decumbens*. *EXCLI Journal* 11: 468 – 479.
- Sasson, A. (1990). Feeding Tomorrow's World. UNESCO Paris: 500-510.
- Sokoto Journal of Veterinary Science (2011) 9 (1):7-11.
- Lasisi, T.K. (2010). Comparative study of body weight gain of savanna brown goat of different age groups raised semi-intensively. A B.Tech project submitted to the Department of Animal Production, School of Agriculture and Agricultural Technology, Federal University of Technology, Minna.
- Ng'ang'a C.J., Maingi N., Kanyari P.W.N., Munyua W.K. (2004). Development, survival and availability of gastrointestinal nematodes of sheep on pastures in a semi-arid area of Kajiado District of Kenya. *Veterinary Resource Communication*. 28(2):491-501
- NSADP (1995). 31st quarterly report, January-March, Niger State Agricultural Development Project Project. pp :32-33.
- Radostitis, O.M., Clive, C.G., Dougins, C.D. and Kenneth, W.H. (2000). A text book of the disease of cattle, sheep, pigs, goats and horses. 9th edition Book power formally ELST with Saunders. Pp 1380-1382.
- Rahamn M.H. (2004). Problems of parasitic diseases among domesticated ruminants in Bangladesh. *The veterinarian*. 12(8): 1 – 4.
- Statistical Analysis System (2008). SAS (STAT9.2) User's Guide. North Carolina, USA: SAS Institute Incorporation.
- Steel, R. G.D. and Torrie, J.H. (1980). Principles and procedures of statistics: A biometric approach, 2nd edition. McGraws Hill Book Company Inc., New York .
- Schneider, S.H. (1989). The Greenhouse Effect: Science and Policy. *Science* 243: 771-781.
- Shakhashiri, B. Z. (2009) Ethanol; Chemical of the week. www.scifun.chem.wisc.edu/chemweek/pdf/ethanol.pdf
- Zaldivar, J., Nielsen, J., and Olsson, L. (2001). Fuel ethanol production from lignocellulose: a challenge for metabolic engineering and process integration. *Applied Microbiology and Biotechnology* 56: 17-34.

HAEMATOLOGICAL AND BIOCHEMICAL INDICES OF JAPANESE QUAILS (*Coturnix coturnix Japonica*) FED GRADED LEVELS OF AFRICAN LOCUST BEAN (*Parkia biglobosa*) PULPAS AN ENERGY SOURCE

¹Yakubu, U. A., ¹Mbahi, T. F., ²Danladi, Y. and ¹Bathon, C. A.

¹Department of Animal Science and Range Management, ModibboAdama University of Technology Yola

²Department of Animal Science, Taraba State University, Jalingo

Corresponding author: uhwesi.yakubu@yahoo.co.uk

Received 8th June, 2016 Accepted 6th September, 2016

Abstract

This study was designed to evaluate the effect of diets containing African Locust bean (*Parkia biglobosa*) pulp on the haematological and biochemical indices of Japanese quails (*Coturnix coturnix japonica*). The study lasted for six (6) weeks during which 240 day old Japanese quails were randomly assigned to Five (5) dietary treatment groups with four (4) replicates in a Completely Randomized Design. The African locust bean pulp was analyzed to contain a crude protein of 6.49%. The birds were fed diets containing graded levels of African Locust Bean pulp at 0, 5, 10, 15 and 20% for T1, T2, T3, T4 and T5 respectively. There was no significant difference ($P>0.05$) observed in the haematological parameters. Biochemical parameters like Creatinine, Urea and Glucose were significantly different ($P<0.05$) across the treatment groups, while Total protein, Albumin, Globulin and Cholesterol were not significantly different ($P>0.05$) across the treatment groups. Therefore, the study revealed that African Locust Bean pulp can be incorporated into the diets of Japanese quails up to 20% as an energy source without any deleterious effect on the blood chemistry of the birds.

Keywords: Japanese quails, African Locust Bean pulp, Haematology, Biochemical Indices

Introduction

The global human population has been on a significant increase. According to the Food and Agriculture Organization of the United Nations (FAO, 2013a; CAST, 2013), demand for food, fuel, and fiber will thus increase to 60% by the year 2050. The population of Nigeria has been on the increase to over 140million (NPC, 2006). This will definitely aggravate the animal protein consumption deficiency of the country. The average animal protein as estimated was put at 7.4g per capita/day as compared with 38g per capita/day of animal protein consumed in South Africa (FAO, 2006). This shows that there is a need to increase the protein intake by intensifying livestock production to a level which can be compared to that of the developed nations.

Poultry occupies a unique position in the livestock sector because they are highly prolific and good converters of feed to meat (Obioha, 1992 and Bot *et al.*, 2012). The domestic fowl has been one of the major sources of animal protein to the Nigerian populace, providing both meat and egg (Odunsi *et al.*, 2007). Emphasis has been on the domestic fowl production whereas nutritive and economic benefits can also be derived from

quail production since quail is fast growing and resistant to many diseases than the domestic fowl (Oluyemi and Roberts, 2000). This shows that there is a problem of pressure on the domestic fowl which can be reduced by widening the scope of meat and egg supply through the population of relatively underutilized poultry species of which Japanese quail is one. Owen and Dike (2013) while reviewing the work of Ani *et al.* (2009) reported that the Japanese quail (*Coturnix coturnix japonica*) was introduced into Nigeria to expand the poultry sub-sector and help supplement the domestic chicken production through meat and eggs.

The largest single dietary need of an animal is for energy. In terms of total cost, energy is the most expensive item in animal diet because of the amount required (Olomu, 2011). Acute shortage and high cost of feed ingredients such as maize, millet and guinea corn have been identified as major hindrances to the expansion of the poultry industry in Nigeria and other developing countries (Fasuyi, 2005 and Bot *et al.*, 2012). Meeting the protein requirement of a developing country like Nigeria will therefore require sourcing for