

INFLUENCE OF UREA AND SHEEP-DROPPINGS ON SOIL pH CHANGES IN SANDY LOAM SOILS OF YOLA, NORTH EASTERN, NIGERIA

Bapetel, U., Tahir, A. M., Bubarai, M. L. and Dadasuko, N.

Department of Soil Science, Modibbo Adama University, Yola, Adamawa state

Corresponding Author's Email Address: usmanubapetel@gmail.com

Received 30th June 2024 Accepted 19th August, 2024

ABSTRACT

An experiment was conducted to assess the influence of urea and sheep dropping applications on soil pH changes in sandy loam soils of Yola, North Eastern Nigeria. The experimental layout was completely randomized design (CRD) with three main factors. The factors were half of recommended rate (50%), full recommended rate (100%) and one and half (150%) recommended rate. The treatments were urea, sheep droppings and combination of urea and sheep droppings. The experiment was replicated three times giving a total of 36 pots. The soils texture was sandy loam, bulk density was below the critical value. The soil was slightly acidic, electrical conductivity was low. The soil had low organic matter and total N, and medium available P levels. Exchangeable Ca was low, Mg and Na were medium and exchangeable K was high. The fertility status of the soils was rated low to medium and could only produce moderate to low yield for most crops. Application of urea, sheep droppings and their combination at half (50%) of the recommended rate, recorded highest pH of 7.40 at 6th week of the experiment on pots amended with 25 urea ha⁻¹ + 2.5 tons SD ha⁻¹. Application of the recommended rate (100%), recorded highest pH value of 7.55 at first week of the experiment on pots amended with 10 tons PD ha⁻¹, while application of one and half (150%) of the recommended rate, recorded highest pH value of 7.85 at 3rd week due to application of PD at the rate of 15 tons ha⁻¹. On the overall the application of SD increased pH from slightly acidic to neutral due to application of 5 tons (50%) and 10 tons (100%) of the recommended rate, while the pH values changed from slightly acidic to slightly alkaline on pots amended with SD at 15 tons ha⁻¹ (150%) of the recommended rate. Based on the results, application of 10 tons SD ha⁻¹ could be recommended for amelioration of soil acidity in sandy loam soils of Yola, North Eastern, Nigeria. Field experiment could also be recommended to ascertain the result obtained.

Key words: urea, sheep droppings, sandy loam soils

INTRODUCTION

Soils in the Nigerian Savannah agro-ecological zones are highly weathered and fragile with low activity clays. The soils are predominantly sandy in texture, low organic matter content, poor water-holding capacity, low nutrient availability as well as buffering capacity (Odunze, 2006). Due to the nature of soil, nutrient depletion is not only common but also constitute a serious constraints to sustainable crop production (Shehuet *al.* 2015). Low pH and lack of appropriate soil fertility restoration practices has also been reported as some of factors that leads to low crop yield in the area (FMARD, 2022).

Soil pH influence nutrients availability, controls biological processes and serves as important parameter for amelioration of problem soils (Jackson and Meetei, 2018). Generally, soil pH is affected by the mineral composition of soil parent material, heavy rainfall, and application of nitrogenous fertilizers (Halilu, *et al.*, 2023). Application of urea and sheep droppings has been reported to

increase crop production in Nigeria and many parts of world (Waniyo, *et al.*, 2013; Ahmet, 2019). The increase in crop yield due to fertilizer application is expected to have positive impact on food production and improvement of livelihood among local farmers (Usman and Kundiri, 2016; Sajidet *al.*, 2016; Mbabahet. *al.*, 2024).

In addition to supply of essential nutrients and increase in crop yield, application of urea and sheep droppings could affect soil pH and availability of some nutrients. Lungu, and Dynoodt, (2008), conducted experiment on the effects of urea application on soil pH and reported that applications of urea resulted in decrease of soil pH, particularly on plots with higher application rates of urea. Decrease in exchangeable Ca and Mg was also observed on the plots treated with urea. This indicates that continuous application of urea could have a negative impact on soil pH and nutrients availability.

On the other hand, application of animal manures could increase soil pH of an

acid soil together with reduction in exchangeable acidity (Ano and Ubochi, 2007). This is an indication that application of urea and organic fertilizers could not only supply essential nutrients but also ameliorates extreme soil pH and moderates other soil properties (Tosin and Jamal, 2012). Thus, this experiment was conducted to assess the effects of urea and sheep droppings on soil pH variations in sandy loam soils of Yola, North East, Nigeria.

MATERIALS AND METHODS

Study Area

The experiment was conducted in the Department of Soil Science Faculty of Agriculture ModibboAdama University Yola, Adamawa State, Nigeria. It is located on longitude 9.35° N and latitude 12.5°E. Mean annual rainfall of the study ranges between 700 to 1000mm, while annual temperature ranges from 15.2 – 39°C. Relative humidity varies seasonally in the state. It is extremely low (20 – 30 %) between January and March. It starts increasing as from April and reaches its peak (above 70 %) in August and September (Adebayo *et al.*, 2020).

The area is characterized by high grassland with many shrubs and few trees. Common grasses in the study area include: *Cassia ssp.*, *Chloris pilosa*, and *Tridax precumbens*. The soil in the study is deep, sandy loam on the surface and sandy clay loam in the sub-surface. The colour of the soil is yellowish brown and poorly drained. The pH is slightly acidic (6.5 – 6.9) with low organic carbon, total N, available P and CEC but medium in ESP and PBS. The soil of the area was classified as Alfisols and Inceptisols (Musa *et al.*, 2007). The geological formation of the soil consists of sandstone, shales with thin limestone at some places (Musa *et al.*, 2007).

EXPERIMENTAL LAYOUT

The experiment design was completely randomised design (CRD) which consisted of three main factors and three treatments and control. The three main factors were (50%) of recommended rate, full dose of recommended rate (100%) and one and half dose (150%) of recommended. For urea, the

treatments were: (T₁) = 50kg, T₂ = 100kg and T₃ = 150kg urea ha⁻¹. For SD, the treatments are: T₁ = 5 tons, T₂ = 10tons and T₃ = 15 tons SD ha⁻¹. While for F₃, the treatments were: T₁ = 25kg urea ha⁻¹ + 2.5 tons SD ha⁻¹, T₂ = 50kg urea ha⁻¹ + 5 tons SD ha⁻¹ and T₃ = 75kg urea ha⁻¹ + 7.5tons SD ha⁻¹. The experiment was replicated three times with the total of 36 pots containing 20kg soil per each pot.

SOIL SAMPLING AND ANALYSIS

Soil samples were taken from all the pots prepared and stored for physical and chemical analysis. Soil texture was determined by the Bouyoucos hydrometer method. Soil pH was measured using pH meter in a 1:2.5 soil-water suspension. Organic carbon and total nitrogen were determined by Walkley-Black oxidation and Micro-Kjeldahl distillation methods respectively. Available P was determined by Bray 1 method. Exchangeable cations were analysed using EDTA (diethylene triaminepenta acetic acid) titrimetric method (FAO, 2020).

RESULTS AND DISCUSSIONS

Soil Physical Properties

Physical properties for soil of the study area is presented in Table 1, the result indicates that sand particles was dominant among the soil separates which had the mean of 67.80%. The sand particles recorded, is lower than 87.83% earlier reported for the soils of the area (Tahir, *et al.*, 2022), though is similar with other reports within northern savannah region of Nigeria (Abdulrashid, *et al.*, 2024). The silt was generally low (mean = 9.87 %). The clay content was medium (mean = 22.33 %) and within the range of 2.8 to 30.2 % earlier reported by Bapetel *et al.* (2024) for soils of Boramji floodplain. The soil textural class was predominantly sandy loam, while the bulk density of the soil was lower than the critical value of 1.65g cm⁻³ but within limit for soils of the region (Ezeaku, *et al.*, 2022). The soil porosity is relatively high indicating the predominance of macro-pores which is a typical characteristic of savannah sandy soils of Nigeria (Moberg, and Esu, 1991).

Table 1: Physical Properties of Soil Used In the Experiment.

Sample	Sand (%)	Silt (%)	Clay (%)	Textural Classes	B.D. (%)	P.D. (%)	Porosity (%)	WHC (%)
T ₁	71.2	8.8	20.0	SL	1.47	2.66	44.74	22.93
T ₂	62.0	9.0	29.0	SCL	1.39	2.67	47.94	32.52
T ₃	70.2	11.8	18.0	SL	1.48	2.65	44.15	23.59
Mean	67.80	9.87	22.33		1.45	2.66	45.61	26.35

Chemical Properties

The soil chemical properties presented in Table 2 indicates that the soil pH of the study area was 6.37 (mean) and rated as slightly acidic (Usman, 2005) and is higher than the mean of 5.7 reported for soils of Boramji Benue River floodplain (Bapetelet *et al.*, 2024), though the pH observed in this study is within the range reported for some soils of Northern savannah region (Jimoh, *et al.*, 2016). Slightly acidic soil pH could not affect nutrient availability since most of the nutrients are released when the soil pH is slightly acidic to neutral in reaction (Jackson, and Meetei, 2018). Electrical conductivity, organic matter and total nitrogen were low, though similar to what was earlier reported (Tahir *et al.*, 2022). Available phosphorus was medium which recorded a mean of 13.33 mg kg⁻¹, but is higher than the mean of 5.80 mg kg⁻¹ earlier reported for Geriyofloodplain (Bapetel, *et al.*, 2021). Higher available P recorded could be attributed to the deposition of silt along with

dissolved particles of P when flood water have receded (Tahir *et al.*, 2006).

The exchangeable Ca of the soil in the area was low (4.61 cmol (+) kg⁻¹), exchangeable Mg was medium (2.34 cmol (+) kg⁻¹), similarly exchangeable Na was also medium (0.52 cmol (+) kg⁻¹), while exchangeable K was high (0.64 cmol (+) kg⁻¹) based on the ratings of Usman (2005) for soils of Adamawa State. The amount of the exchangeable bases is similarly to what was reported for the soils of the area (Alpha *et al.*, 2011). The soils also had low exchangeable acidity and high base saturation but is similar with what was earlier reported by Moberg and Esu, (1991). With low organic matter content, total nitrogen and exchangeable bases, soil fertility status for the soils could be rated as low and it agrees with earlier results for the study area (Bapetel *et al.*, 2024). Soils with low fertility status could only support low yield and therefore require fertilizer application for sustainable cultivation of crops

Table 2: Chemical Properties for Soil of the Study Area.

Sample	pH	EC	OC	OM	TN	AvP	Ca	Mg	Na	K	TEB	TEA	ECEC	PBS
	(1:2.5)	(dS m ⁻¹)				(mg kg ⁻¹)								(%)
										Cmol kg ⁻¹				
T ₁	6.27	0.15	1.08	1.86	0.11	13.51	4.61	2.11	0.41	0.621	7.751	2.25	10.00	77.50
T ₂	6.48	0.19	0.92	1.58	0.09	11.21	5.11	2.01	0.62	0.632	8.372	1.98	10.35	80.87
T ₃	6.36	0.21	0.81	1.39	0.08	15.28	4.11	2.89	0.53	0.67	8.20	2.31	10.51	78.02
Mean	6.37	0.18	0.94	1.61	0.09	13.33	4.61	2.34	0.52	0.64	8.11	2.18	10.29	78.80

EC = Electrical conductivity, OC = Organic carbon, OM = Organic matter, TN = Total nitrogen, Av P = Available phosphorus, Ca = Calcium, Mg = Magnesium, K = Potassium Na = Sodium, TEB = Total exchangeable bases, TEA = Total exchangeable acidity, ECEC = Effective cation exchange capacity, PBS = Percentage base saturation

Half (50%) of the Recommended Rates of Urea and Sheep droppings (SD)

Effects of the application of half dose (50%) of the recommended rates of urea and poultry droppings on soil pH (Table 3) indicated that there were significant effects among the treatments except at 2nd and 4th weeks after application. During the first week of the experiment, highest pH of 6.26 was recorded on pot amended with 5 tons ha⁻¹ of sheep

droppings (SD), at second week, highest pH of 6.25 was recorded on pots amended with urea and SD at the rates of 25 kg and 2.5 tons ha⁻¹ were applied. For week three, application of 5 tons SD ha⁻¹ gave the highest pH of 7.21. At 5th week of the experiment, highest pH of 6.43 was recorded on pots amended with 5 tons SD ha⁻¹, while at the 6th week highest pH of 7.40 was recorded on pots where 25 kg urea and 2.5 tons SD ha⁻¹ were applied.

Highest pH value of 7.40 was recorded on pots amended with 25kg urea + 2.5tons SD ha⁻¹, followed by pH of 7.21 recorded at the third week of the experiment on pots amended with 5 tons SD ha⁻¹ (Table 3). The highest pH of 7.40 recorded is within the neutral range (Usman, 2005) but is higher than what was reported for the soils of the region (Haliluet *al.*, 2023). The result indicates that the soil pH changed from slightly acidic on the control plots to neutral with the application

of SD alone as well as combined application of urea and SD. The observation agrees with what was earlier reported by Ano and Ubochi (2007). Neutral soil pH is more conducive for maximum nutrient release and availability, which also favours increased microbial activities along with release of nutrients embedded in organic substances (Jackson and Meetei, 2018).

Table 3: Changes of Soil pH due to 50% below the Application rates of Urea and Sheep Droppings

TRT/Weeks	W1	W2	W3	W4	W5	W6
Control	5.41 _c	5.84 ^a	6.61 ^b	6.20 ^a	5.66 ^d	6.30 ^b
50 kg urea ha ⁻¹	5.78 ^b	6.20 ^a	6.28 ^c	6.25 ^a	6.43 ^a	5.65 ^b
5 tons SD ha ⁻¹	6.26 ^a	5.75 ^a	7.21 ^a	6.20 ^a	5.91 ^c	5.67 ^b
25 kg urea ha ⁻¹ + 2.5 tons SD ha ⁻¹	6.20 ^a	6.25 ^a	6.32 ^c	6.33 ^a	6.22 ^b	7.40 ^a
SE	0.13	0.55	0.066	0.58	0.049	0.45

TRT = Treatment, WK1WK6 = week 1 to week 6, SE = Standard error, SD = Sheep droppings

Full (100%) of the Recommended Rates of Urea and Sheep droppings (SD)

Effects of the application of recommended rates (100%) of urea, SD and their combination is presented in Table 4. The results indicated that there were significant effects of the application throughout the six weeks of the experiment. For the first week, highest pH value of 7.55 was recorded on pots amended with 10 tons SD ha⁻¹, at second week highest pH of 7.45 was recorded on pots that received combined applications of urea and SD at the rates of 50kg urea + 5tons SD ha⁻¹. At three weeks, there was a slight decrease in soil pH, with the highest of 6.33 recorded on pots amended with 10 tons SD ha⁻¹, at 4th week of the experiment, highest pH of 7.15 was also recorded on pots amended with 10 tons SD ha⁻¹. General decrease in soil pH at 5th and 6th weeks of the experiment were observed, with the highest pH values of 6.34 and 6.45 recorded on pots amended with 10 tons SD ha⁻¹.

Overall highest pH recorded during the first week of experiment in pots amended with 10 tons SD ha⁻¹ is within the neutral range, though higher than what was earlier reported for study area and other areas within savannah agro-ecological zone (Shehuet *al.*, 2015; Tahir *et al.*, 2022; Ezeaku, *et al.*, 2022). Highest pH value recorded during the first week of the experiment could be attributed to the increase in pH due to nutrient released from product of decomposition of sheep droppings. Alexandra and Jose (2015) reported that when organic amendments decomposed, they are converted into CO₂ and humus with a simultaneous release of nutrients that could increase the soil pH. According to Ano, and Ubochi (2007), animal manures significantly increased soil pH from 4.6 to 5.6. These are good indications that application of urea and organic fertilizers could not only supply essential nutrients, but also ameliorates extreme soil pH and moderates other soil properties such as water-holding capacity (Tosin and Jamal, 2012).

Table 4: Changes in Soil pH due to Application of the Recommended Rate of Urea and Sheep Droppings

TRT/Weeks	W1	W2	W3	W4	W5	W6
Control	5.41 ^d	5.41 ^b	5.61 ^a	5.20 ^a	5.66 ^b	5.30 ^a
100kg/ha Urea	5.65 ^c	6.22 ^{ab}	6.33 ^b	6.30 ^a	5.66 ^b	6.20 ^{ab}
10 tons/ha SD	7.55 ^a	5.60 ^b	6.28 ^b	7.15 ^a	6.34 ^a	6.45 ^a
50kg/ha U+5tons/ha SD	6.21 ^b	7.45 ^a	6.26 ^b	6.20 ^a	5.37 ^c	5.92 ^b
SE	0.082	0.628	0.059	0.585	0.067	0.153

TRT = Treatment, WK1WK6 = week 1 to week 6, SE = Standard error, SD = Sheep droppings

One and Half (150%) of the Recommended Rates of Urea and Sheep droppings (SD)

Effects of the application one and half recommended rates for Urea, SD and their combinations is presented in Table 5. There were significant effects of the application except at 4th week of experiment where no significant effect was recorded. At first week, highest pH of 5.91 was recorded due to application of 15 tons SD ha⁻¹. At second week, application of both 150 kg urea and 15 tons SD ha⁻¹ gave the highest pH of 5.90. At the third week of the experiment, highest pH of 7.85 was recorded on pots where 15 tons SD ha⁻¹ was applied. Similarly, the highest pH

of 6.27 was recorded at 5th week with the application of 15 tons SD ha⁻¹, while highest pH of 6.33 was recorded at 6th week of the experiment on pots where combined applications of urea and SD at the rate of 75 kg urea ha⁻¹ + 7.5 tons SD ha⁻¹ were made. Highest pH value of 7.85 recorded at the third week of the experiment is slightly alkaline in reaction based on ratings for the soils of the area (Usman, 2005). Alkaline soil pH could benefit microbial activities and availability of some essential nutrients, but it could have some detrimental effects on the availability of micronutrients (Jackson and Meetei, 2018; Israel and Mira, 2021).

Table: 5 Changes in Soil pH due to the Application of 150% above the Recommended Rate of Urea and Sheep Droppings

TRT/Weeks	W1	W2	W3	W4	W5	W6
Control	5.41 ^d	5.41 ^b	5.61 ^c	5.20 ^a	5.66 ^b	5.30 ^a
150kg/ha Urea	5.91 ^{ab}	5.90 ^a	6.83 ^b	6.32 ^a	5.53 ^b	5.30 ^b
15 tons/ha SD	5.34 ^a	5.90 ^a	7.85 ^a	6.15 ^a	6.27 ^a	5.65 ^b
75kg/ha U+7.5tons/ha SD	5.19 ^a	6.35 ^a	6.28 ^d	6.50 ^a	5.62 ^b	6.33 ^a
SE	0.24	0.68	0.059	0.46	0.064	0.18

TRT = Treatment, WK1WK6 = week 1 to week 6, SE = Standard error, SD = Sheep droppings

CONCLUSION

The soil of the study area was slightly acidic with the mean of 6.37, electrical conductivity was low (0.18). The soil had low organic matter, total nitrogen and medium P levels. Exchangeable Ca was low, exchangeable Mg and Na were medium and exchangeable K was high. The fertility status for the soils was rated low to medium and could produce moderate to low yield for most crops.

On the overall the application of SD increased soil pH from slightly acidic to neutral due to application of 5 tons (50%) and 10 tons ha⁻¹ (100%) of the recommended rate, while the pH values changed from slightly acidic to slightly alkaline due to application of SD at 15 tons ha⁻¹ (150%) of the recommended rate. Based on the results, application of at 10 tons SD ha⁻¹ could be recommended for amelioration of acidity in sandy loam soils of Adamawa State. Field experiment could also be recommended to ascertain the result obtained.

REFERENCES

- Abdulrashid I., Adeduntan, S. A. Adekunle, V.A.J., and Wali B. R. (2024). Assessment of Soil Physicochemical Properties in the Parklands of Northern Nigeria. *African Journal of Environment and Natural Science Research*, 7(1):146-154.
- Adebayo, A. A., and A. L. Tukur, A. A. Zemba (2020). Adamawa State in Maps. 2nd Edition, Paraclete Pub Yola, Nigeria. 203pp
- Ahmet, P. (2019). The Effect Of Sheep Manure On Yield And Quality In Production Of Organic Sugar Beet. *Israel Journal of Plant Sciences*, <https://www.researchgate.net/publication/33546915>. 5p
- Alexandra, B. and Jose, B. (2015). The importance of soil organic matter: Key to drought-resistant soil and sustained food production. Food and Agriculture Organization of The United Nations, Rome.
- Alpha Y. K., Friday E., Lucky O. O. and Hakeem A. A. (2011). Phosphorus and Nitrogen Fertilization of

- Soybean in the Nigerian Savanna. Explain Agriculture, Cambridge University Press: pp: 1 – 10.
- A. M. Tahir, A. M. Saddiq and B. H. Usman. (2006). Effect of Seasonal Flooding on Particle Size Sedimentation on Soils of Geriyo Irrigation Project (GIP), Yola, Nigeria. *Nigerian Journal of Tropical Agriculture*. Vol. 8: 120 – 129.
- Bapetel, U., Tahir, A. M., Bubarai M. L, and Michael, J. A. (2024). Assessment of Soil Fertility Status of Boramji Irrigation Project Site, Yola North, Adamawa State, North-East Nigeria. *Nigerian Journal of Tropical Agriculture*, 28: 207 – 215.
- Bapetel, U., Isa, S. S., Tahir A. M. and Abdulkadir, N. A. (2021). Assessment of Fertility Status of Soil of Geriyo Irrigation Project, Adamawa State, Nigeria. *Nigerian Journal of Tropical Agriculture*. 21: 102 – 111.
- Ezeaku, P.I., Kefas, P.K., Hashim, S.A., Ali, I.J., Gisilanbe, S.A., Kussy, J.J., Garpiya, B.T., Yusuf, B., Udoh, S., Ezeaku, V.I. and M.A.N Anikwe. (2022). Characterization, Taxonomic and Fertility Capability Classification of some Soils in the Teaching and Research Farm of Taraba State University, Jalingo. *Nigerian Journal of Soil Science*. 32(1): 77 – 82.
- FMARD, (Federal Ministry of Agriculture and Rural Development). (2022). National Agricultural Soil Management Policy (Final Draft Copy). 71pp
- FAO. (2021). Standard operating procedure for soil pH determination Standard operating procedure for soil pH determination. Rome. 23p
- Halilu S., Omomoh E., Nannim S., Gujahir R., Danlami R., Moses O. O. and Gwamzhi E. P. (2023). Soil pH Variability in Mangu, Bokkos and Pankshin Local Government Areas of Plateau State. *World Journal of Advanced Research and Reviews*. 19(03): 1508 – 1517
- Israel, Z. and Mira, S. (2021). The effects of alkalinity on physical and chemical properties of soil. *PULJPBAS*, 3(2):1 – 5.
- Jackson, K. and Meetei, T. T. (2018). Influence of soil pH on nutrient availability: A Review. *Journal of Emerging Technologies and Innovative Research (JETIR)* www.jetir.org. 5 (12):708 – 713.
- Jimoh, A.I., Malgwi, W.B., Aliyu, J. and Shobayo, A.B. (2016). Characterization, Classification and Agricultural Potentials of Soils of Gabari District, Zaria, Northern Guinea Savanna Zone Nigeria. *Biological and Environmental Sciences Journal for the Tropics*. 13(2) September, 13(2): 102 – 113.
- Lungu, O.I.M. and Dynoodt, R.F.P. (2008). Acidification from Long-Term Use of Urea and Its Effect on Selected Soil Properties Mbabah, S. N., Anyanwu, N. J., Idiong, G. C., Eketete, I. E. and Johnson, P. E. (2024). Impact of Different Manure on the Growth and Yield of Maize in Southern Nigeria. *Animal Research International*, 21(1): 5257 – 5264.
- Moberg, J.P. and Esu, I. E. 1991. Characteristics and Composition of Some Savanna Soils in Nigeria. *Geoderma*. 48(1-2):113 – 129.
- Musa, H., Singh, L., Usman, B.H., Saddiq, A.M. and Mustapha, S. (2007). Characterization and Classification of the soil developed on basement complex in Federal University of Technology, Yola, Nigeria. *JOLORN*, 8:183 – 189.
- O. Ano and C. I. Ubochi. (2007). Neutralization Of Soil Acidity By Animal Manures: Mechanism of Reaction. *African Journal of Biotechnology*. 6 (4): 364 – 368.
- Odunze, A.E.(2006). Soil properties and management strategies for some sub-humid of savanna zone Alfisols in Kaduna State, Nigeria. *Samaru J. Agric. Res.*, 22: 3 – 14.
- Sajid K., Mohammad T., Bilal A., Tariq S. and Wisal A. (2016). Yield and Yield

- Components of Maize As Affected By Integrated Management of Sheep Manure and Urea Fertilizer. *Middle East Journal of Agriculture Research*. 5 (4): 522 – 533.
- Shehu, B.M. Jibrin, J.M. and Samndi, A.M. (2015). Fertility Status of Selected Soils in the Sudan Savanna Biome of Northern Nigeria. *International Journal of Soil Science* 10 (2): 74 – 83.
- Tahir, A. M., Bapetel, U., Bubarai, M. L. And Leyiro, A. (2022). Fertility Status For Soils Of Teaching And Research Farm, Modibbo Adama University, Yola, Sudan Savanna Zone Of Nigeria. *Taraba Journal of Agricultural Research*, 10 (1): 24 – 32.
- Tosin A. G. and Jamal O. A. (2012). Effects Of Farm Practices And Manure Application On Soil Physicochemical Properties And Proceeds Of Maize Grown On A Degraded Intensively Tilled Alfisol In Southwestern Nigeria. *International Journal of Manures and Fertilizers*. 1 (8): 144 – 154.
- Usman, S. and Kundiri, A. M. (2016). Values of Organic Materials as Fertilizers to Northern Nigerian Crop Production Systems. *Soil and Environmental Research*. 7(12): 204 – 211.
- Usman, B.H. (2005). The soils of Adamawa State, North-eastern Nigeria. In: Igwe, E.C., Mshelia, S.I. and Jada, M.Y. (eds). *Agriculture in Adamawa State*. pp 62-83.
- Waniyo, U.U., Sauwa, M.M., Ngala, A.L., Abubakar, G.A. and Anelo, E.C. (2013). Influence of Sources and Rates of Manure on Yield and Nutrient Uptake of Maize (*Zea mays* L.) in Maiduguri, Nigeria. *Nigerian Journal of Basic and Applied Science*, 21(4): 259 – 265.